arXiv:hep-ph/0502043v1 4 Feb 2005

Can there be neutrino oscillation in Gamma-Ray Bursts fireball ?

Sarira Sahu and Juan Carlos D'Olivo
Instituto de Ciencias Nucleares,
Universidad Nacional Autónoma de México,
Circuito Exterior, C.U., A. Postal 70-543,
04510 Mexico DF, Mexico

The central engine which powers the Gamma-Ray Burst (GRB) fireball, produces neutrinos in the energy range of about 5-20 MeV. Fractions of these neutrinos may propagate through the fireball which is far away from the central engine. We have studied the propagation of these neutrinos through the fireball which is contaminated by baryons and have shown that, resonant conversion of neutrinos are possible for the oscillations of $\nu_e \leftrightarrow \nu_{\mu,\tau}$, $\nu_e \leftrightarrow \nu_s$ and $\bar{\nu}_{\mu,\tau} \leftrightarrow \bar{\nu}_s$ if the neutrino mass square difference and mixing angle are in the atmospheric and/or LSND range. On the other hand it is probably difficult for neutrinos to have resonant oscillation if the neutrino parameters are in the solar neutrino range. From the resonance condition we have estimated the fireball temperature and the baryon load in it.

PACS numbers: 14.60.Pq,98.70.Rz

GRBs are short, non-thermal bursts of low energy ($\sim$ 100 KeV-1 MeV) photons and release about 10^{51}-10^{53} ergs in a few seconds making them the most luminous object in the universe. A class of models call *fireball model* seems to explain the temporal structure of the bursts and the non-thermal nature of their spectra[1, 2, 3, 4, 5]. Sudden release of copious amount of γ rays into a compact region with a size $c\delta t \sim$ 100-1000 Km[1] creates an opaque $\gamma - e^{\pm}$ fireball due to the process $\gamma + \gamma \rightarrow e^+ + e^-$. The average optical depth of this process[6] is $\tau_{\gamma\gamma} \simeq 10^{13}$. This optical depth is very large and even if there are no pairs to begin with, they will form very rapidly and will Compton scatter lower energy photons. Because of the huge optical depth, photons can not escape freely. In the fireball the γ and $e^{\pm}$ pairs will thermalize with a temperature of about 3-10 MeV. The fireball expands relativistically with a Lorentz factor $\Gamma \sim 100 - 1000$ under its own pressure and cools adiabatically due to the expansion. The radiation emerges freely to the inter galactic medium (ISM), when the optical depth is $\tau_{\gamma\gamma} \simeq 1$.

In addition to γ, $e^{\pm}$ pairs, fireballs may also contain some baryons, both from the progenitor and the surrounding medium. These baryons can be either free or in the form of nuclei. If the fireball temperature is high enough (more than 0.7 MeV), then it will be mostly in the form of neutrons and protons. Derishev et al. [7, 8] argued that roughly equal numbers of neutrons and protons should be present in the fireball. The electrons associated with the matter (baryons) can increase the opacity, hence delaying the process of emission of radiation and the baryons can be accelerated along with the fireball and convert part of the radiation energy into bulk kinetic energy. But irrespective of it, the baryonic load has to be very small, otherwise, the expansion of the fireball will be Newtonian, which is inconsistent with the present observations. Why the baryonic loading is so low in the fireball is still an open question to be answered[1]. The neutrino oscillation may overcome the baryon loading problem[9, 10] ! The evolution of the pure fireball (with no baryons) has been studied in ref.[11, 12]. In the expanding fireball, protons are accelerated in shocks and collide with the photons to produce charged pions, which give rise to ultra high-energy neutrinos[13]. These neutrinos can be detected by km^2 detectors. Observation of these neutrinos can study oscillating neutrino flavors in the largest possible baseline, test the equivalence principle and many other neutrino properties [14].

The hidden central engine which powers the fireball is still unknown, but observation suggests that it must be compact. The prime candidates are merger of neutron star with neutron star (NS-NS), black hole-neutron star binaries (BH-NS), hypernova/collapsar models involving a massive stellar progenitor[1, 2, 3, 4, 15]. The recent observations of GRB030329 strongly favor collapsar model [16]. In all these models, the gravitational energy is released mostly in the form of $\nu\bar{\nu}$, gravitational radiation and a small fraction ($\sim 10^{-3}$) is responsible to power GRB. Neutrinos of energy about 5-20 MeV are generated due to the stellar collapse or merger event that trigger the burst. Also due to nucleonic bremsstrahlung $NN \rightarrow NN\nu\bar{\nu}$ and $e^+e^- \rightarrow \nu\bar{\nu}$ processes muon and tau types neutrinos can be produced[17] during the merger process and its flux will be very small. Fractions of these neutrinos may propagate through the fireball which is far away from the central engine. Also within the fireball, because of the weak interaction process $p + e^- \rightarrow n + \nu_e$, MeV neutrinos can be generated and propagate through it. The fireball plasma being in an extreme condition, may affect the propagation of MeV neutrinos through it.

In a heat bath the dispersion relation which governs the propagation of the particle gets modified and this can have a drastic effect on the particle propagation in the heat bath. The neutrino propagation in a thermal bath has been studied extensively[18, 19, 20, 21]. In general, the propagating neutrino will experience an effective potential due to the particles in the heat bath. Particu-

larly the neutrino propagation in the early universe hot plasma, as well as in the supernova medium has profound implications in the respective physics. For example the neutrino oscillation in the early universe hot plasma may change the relative abundances of ν_e and $\bar{\nu}_e$ and affect the primordial nucleosynthesis of light elements. In the supernova case, the neutrino oscillation in the dense and compact environment can affect the cooling mechanism.

The electron type neutrinos have charge current as well as neutral current interactions, but the muon and tau types will experience only the neutral current interactions. To the leading order the effective potential experience by the neutrinos is proportional to the difference of the particle anti-particle number densities. So if the system under consideration has equal number of particles and anti-particles, the leading order contribution vanishes. On the other hand, the next-to-leading order contribution (i.e. the term proportional to $1/M^4$, where M is the vector boson mass) is proportional to the sum of the particle and anti-particle number densities and this is the leading contribution to the effective potential. The early universe hot plasma, supposed to have almost equal population of particles and anti-particles, hence the leading contribution to the neutrino effective potential will be proportional to $1/M^4$. Similar situation can arise for neutrinos propagating in the GRB fireball if one consider mostly photon-lepton fireball[1, 3, 4]. Here in the present work we consider a fireball containing mostly photon-lepton with little baryon contamination, which mimics the early universe hot plasma and study the neutrino propagation within it.

In a relativistic and nondegenerate $e^{\pm}$, proton and neutron plasma, the effective potential experiences by ν_e is [19, 20]

$$V_{\nu_e} \simeq \sqrt{2} G_F N_\gamma \left[\mathcal{L}_e - \left(\frac{7\xi(4)}{\xi(3)} \right)^2 \frac{T^2}{M_W^2} \right], \tag{1}$$

and by a muon or tau neutrino it is given by

$$V_{\nu_{\mu,\tau}} \simeq \sqrt{2} G_F N_\gamma \mathcal{L}_{\mu,\tau}, \tag{2}$$

where

$$\mathcal{L}_e = \left(\frac{1}{2} + 2\sin^2\theta_W \right) L_e + \left(\frac{1}{2} - 2\sin^2\theta_W \right) L_p - \frac{L_n}{2}, \tag{3}$$

and

$$\mathcal{L}_{\mu,\tau} = \left(-\frac{1}{2} + 2\sin^2\theta_W \right) (L_e - L_p) - \frac{L_n}{2}. \tag{4}$$

The asymmetry of the particle a is defined as

$$L_a = \frac{(N_a - \bar{N}_a)}{N_\gamma}, \tag{5}$$

where $N_\gamma = \frac{2}{\pi^2}\xi(3)T^3$ is the number density of photons. For the anti-neutrinos, the effective potential will be given by changing $L_a \to -L_a$. For a pure γ and $e^{\pm}$ fireball, we have $L_p = L_n = 0$.

We consider the neutrino oscillation processes: $\nu_e \leftrightarrow \nu_{\mu,\tau}$, $\nu_e \leftrightarrow \nu_s$ and the antineutrino processes. The effective potential difference for $\nu_e \leftrightarrow \nu_{\mu,\tau}$ and $\bar{\nu}_e \leftrightarrow \bar{\nu}_{\mu,\tau}$ processes, is given by

$$V \simeq 4.02 \times 10^{-12} T_{MeV}^3 \left[\pm L_e - 6.14 \times 10^{-9} T_{MeV}^2 \right] \ MeV, \tag{6}$$

where $\pm$ corresponds to ν and $\bar{\nu}$ respectively, and for $\nu_e \leftrightarrow \nu_s$ oscillation, the effective potential difference is given by Eq.(1). For pure γ, $e^{\pm}$ fireball (CP-symmetric), $L_e = 0$ and only the higher order term will contribute. The conversion probability for the above processes for a constant V is given by,

$$\mathcal{P}(t) = \frac{\Delta^2 \sin^2 2\theta}{\omega^2} \sin^2(\frac{\omega t}{2}), \tag{7}$$

where

$$\omega = \sqrt{(V - \Delta\cos 2\theta)^2 + \Delta^2 \sin^2 2\theta}, \tag{8}$$

with $\Delta = \Delta m^2/2E_\nu$, V is the potential difference, E_ν is the neutrino energy and θ is the neutrino mixing angle. The oscillation length is given by

$$L_{osc} = \frac{L_v}{\sqrt{\cos^2 2\theta (1 - \frac{V}{\Delta\cos 2\theta})^2 + \sin^2 2\theta}}, \tag{9}$$

where $L_v = 2\pi/\Delta$ is the vacuum oscillation length which can be recovered for $V = 0$. For resonance to occur we should have from Eq.(8)

$$V = \Delta \cos 2\theta. \tag{10}$$

We have so far assumed that lepton asymmetry in the fireball does not vary with distance. But in reality, the lepton asymmetry changes with distance. So in this case we will consider the adiabaticity of the resonant conversion. The adiabaticity condition at the resonance[22] is given by

$$\kappa_r \equiv 2.0 \times 10^{-3} \left(\frac{\tilde{\Delta m^2} \sin 2\theta}{E_{MeV}} \right)^2 \frac{l_{cm}}{T_{MeV}^3} \left(\frac{dL_e}{dx} \right)^{-1} \geq 1, \tag{11}$$

where l_{cm} is some length scale expressed in centimeter, x is a dimensionless variable, $\tilde{\Delta m^2}$ is in unit of eV^2 and T_{MeV} and E_{MeV} are expressed in unit of MeV. The above condition depends on the neutrino parameters and the change in the L_e as one goes away from the center of the fireball.

As already stated in the introduction, baryon loading is still an outstanding problem. It is believed that the contamination is very low ($10^{-8}M_\odot - 10^{-5}M_\odot$)[1, 2], so that, the fireball can have a ultra-relativistic expansion. For simplicity, we consider a charge neutral spherical fireball ($L_e = L_p$) of initial radius R with equal number of

protons and neutrons in it. Then the baryon load in the fireball is

$$M_b \sim 2.23 \times 10^{-4} R_7^3 T_{MeV}^3 L_e M_\odot, \tag{12}$$

where R_7 is in unit of 10^7 cm. As stated above, the baryon contamination is of order $10^{-8} M_\odot$ to $10^{-5} M_\odot$, which corresponds to $L_e \sim 4.47 \times 10^{-5} R_7^{-3} T_{MeV}^{-3}$ to $L_e \sim 4.47 \times 10^{-2} R_7^{-3} T_{MeV}^{-3}$ respectively. The thermalized fireball has a temperature $T \sim (L/4\pi\sigma R^2)^{1/4} \sim$ 3-10 MeV, with L the luminosity and σ the Stephen-Boltzmann constant.

The processes (active to active oscillations) do not depend on the baryon asymmetry in the fireball, simply because, the neutral current contribution to the potential of all the active neutrinos is the same and for active to active oscillation this contribution cancels out, leaving only the dependence on L_e. But the active-sterile oscillation depends on both the lepton and the baryon asymmetry. In Eq.(6), $V > 0$ for neutrino process if

$$L_e > 6.14 \times 10^{-9} T_{MeV}^2, \tag{13}$$

and depending on the fireball parameters and the neutrino properties the resonance condition can be satisfied. On the other hand, antineutrino process will never satisfy the resonance condition because the potential is always negative. The resonance condition of Eq.(10) can be written as

$$L_e T_{MeV}^3 = 0.124 \frac{\tilde{\Delta m}^2 \cos 2\theta}{E_{MeV}}, \tag{14}$$

and the resonance length is given by

$$L_{res} \simeq 248 \frac{E_{MeV}}{\tilde{\Delta m}^2 \sin 2\theta}\ cm. \tag{15}$$

Putting the value of L_e from Eq.(14) in Eq.(13), the constraint on the fireball temperature is

$$T_{MeV}^5 < 0.2 \times 10^8 \frac{\tilde{\Delta m}^2 \cos 2\theta}{E_{MeV}}. \tag{16}$$

Thus the fireball temperature derived in Eq.(16) is the one required in order to have resonant conversion of the neutrinos. As we have already discussed, during the stellar collapse or merger events that trigger the burst, neutrinos of energy about 5-20 MeV are copiously produced and some of these neutrinos will propagate through the fireball. Apart from this, due to inverse beta decay, MeV neutrinos can also be generated within the fireball. So here we will take $E_{MeV} = 5$ and 20 for our calculation to estimate the fireball parameters.

Let us study the resonance condition for Solar, Atmospheric and the reactor (LSND) neutrinos, where we know approximately the neutrino mass square differences and the neutrino mixing angles from the recent experimental results. These can constraint the fireball parameters.

The recent analysis of the salt phase data of SNO[23] combines with the KamLAND[24] reactor antineutrino results gives $6 \times 10^{-5}\ eV^2 < \Delta m^2 < 10^{-4}\ eV^2$ and $0.8 < \sin 2\theta < 0.98$ with a confidence level of 99%. The best fit point has $\Delta m^2 \sim 7.1 \times 10^{-5}\ \ eV^2$ and $\sin 2\theta \sim 0.83$. Using the best fit point in Eq.(14) we obtain $L_e \simeq 0.5 \times 10^{-5} T_{MeV}^{-3} E_{MeV}^{-1}$. The condition in Eq.(16) gives $T_{MeV} < 2.8$ and < 2.1 respectively for $E_{MeV} = 5$ and 20. Similarly the resonance length $L_{res} \sim$ 211 kilometers and 845 kilometers are obtained respectively for $E_{MeV} = 5$ and 20. Using the resonance value of L_e in Eq.(12) we obtain $M_b \sim 1.15 \times 10^{-9} R_7^3 M_\odot E_{MeV}^{-1}$ which is independent of the fireball temperature. For $E_{MeV} = 5$ and 20 we obtain $M_b \sim 0.23 \times 10^{-9} R_7^3 M_\odot$ and $M_b \sim 0.58 \times 10^{-10} R_7^3 M_\odot$ respectively. One can see that, in this case the temperature of the fireball is less compared to the lower limit of $3MeV$. Also the baryon load in the fireball is less, which can be improved by increasing R_7. By considering the solar neutrino mixing angle and the Δm^2, for neutrino oscillation in the fireball, the fireball temperature has to be less than 3 MeV. If the L_e vary with distance then at the resonance $\kappa_r \sim 0.8 \times 10^{-14} l_{cm} (L_e')^{-1}$, which implies, $L_e' = dL_e/dx$ has to be very small to satisfy the condition in Eq.(11).

The Super-Kamiokande Collaboration recently reported the atmospheric neutrino oscillation parameters[25] $1.9 \times 10^{-3}\ eV^2 < \Delta m^2 < 3.0 \times 10^{-3}\ eV^2$ and $0.9 \leq \sin^2 2\theta \leq 1.0$ with a 90% confidence level. Taking $\Delta m^2 \sim 2.5 \times 10^{-3}\ \ eV^2$ and $\sin^2 2\theta \sim 0.9$ we obtain, $L_e \simeq 0.98 \times 10^{-4} T_{MeV}^{-3} E_{MeV}^{-1}$ and this implies $T_{MeV} < 5$ and < 3.8 respectively for $E_{MeV} = 5$ and 20. So these temperature come within the range of the fireball as discussed earlier. The resonance length for $E_{MeV} = 5$ and 20 are respectively 5.2 kilometers and 21 kilometers. Also for the above ranges of neutrino energy, the baryon load in the fireball is $M_b \sim 0.44 \times 10^{-8} R_7^3 M_\odot$ and $M_b \sim 0.11 \times 10^{-8} R_7^3 M_\odot$ respectively. Thus before coming out of the fireball, the neutrinos can have many resonant oscillations. If the lepton asymmetry vary with distance, then $l_{cm}/L_e' \geq 10^{12}$ to have resonant conversion.

Thirdly let us discuss the implication of the LSND and KARMEN results[26] on the neutrino oscillation in the GRB fireball. The combined analysis of LSND and KARMEN 2 results give $0.45\ eV^2 < \Delta m^2 < 1\ eV^2$ and $2 \times 10^{-3} < \sin^2 2\theta < 7 \times 10^{-3}$ with a confidence level of 90%. We consider $\Delta m^2 \sim 0.5\ \ eV^2$ and $\sin 2\theta \sim 0.07$ to estimate the fireball parameters. This gives, $T_{MeV} < 18$ and < 14 respectively for $E_{MeV} = 5$ and 20 and for these two values of E_{MeV} the baryon load is respectively $M_b \sim 0.3 \times 10^{-5} R_7^3 M_\odot$ and $M_b \sim 0.7 \times 10^{-6} R_7^3 M_\odot$. The resonance lengths for these two neutrino energies are respectively 0.4 kilometer and 1.4 kilometer. So this shows that, the propagating neutrinos have to oscillate several times, before they come out of the fireball. If the lepton asymmetry has a profile then in this case $l_{cm}/L_e' \geq 10^{10}$ to satisfy the resonant oscillation.

If the neutrino oscillation parameters are in the at-

mospheric and/or LSND range, the oscillation length at resonance within the fireball will vary from few meters to $\sim$ 21 kilometers. As the resonance length is small compared to the size of the fireball, there will be many resonant oscillations for $\nu_e \leftrightarrow \nu_{\mu,\tau}$ before they emerge out of the fireball. So the average conversation probability in this case is $\mathcal{P}(t) \sim 0.5$, where we have considered the fact that L_e does not vary with distance. But if the oscillation parameters are in the solar neutrino range, probably it is difficult to have neutrino oscillation. This is due to the fact that, the fireball temperature in this case is less (i.e. < 3 MeV).

For the oscillation $\nu_e \leftrightarrow \nu_s$, with charge neutral fireball and $L_p = L_n$, the effective potential is given by

$$V \simeq 4.02 \times 10^{-12} T_{MeV}^3 \left[\frac{L_e}{2} - 6.14 \times 10^{-9} T_{MeV}^2 \right] MeV. \tag{17}$$

Resonance condition can also be satisfied for the above process for $V > 0$. Both the processes $\nu_e \leftrightarrow \nu_{\mu,\tau}$ and $\nu_e \leftrightarrow \nu_s$ are equally probable if we consider the atmospheric neutrino oscillation parameter or the LSND one, with average probability ~ 0.5. Finally let us consider the process $\bar{\nu}_{\mu,\tau} \leftrightarrow \bar{\nu}_s$, for which

$$V \simeq 2.0 \times 10^{-12} T_{MeV}^3 L_e \ MeV. \tag{18}$$

Here the higher order contribution to the neutrino potential is absent and due to this, there is no constraint on the baryon loading of the fireball. Only the charge neutrality of the fireball is sufficient enough for the resonant oscillation to take place.

From the collapsar or merger model of SN with SN and/or from the collision of SN with a black-hole, lots of neutrinos will be produced and fractions of these neutrinos will propagate through the fireball. Here we have studied the propagation of these neutrinos through the GRB fireball by assuming the later to be spherical with a radius R, charge neutral and $L_p = L_n$. Also we have assumed that $L_e > 6.14 \times 10^{-9} T_{MeV}^2$ so that the potential difference will be positive for neutrinos and there can be resonant oscillation. By using the known neutrino mass square difference and mixing angle from the solar, atmospheric and reactor experiments we estimate the fireball temperature, the baryon load and the lepton asymmetry in it. Our result shows that, if the neutrino oscillation parameters are in the solar neutrino range, probably, there will be very few or no oscillation will take place. On the other hand if the neutrino oscillation parameters are in the atmospheric and/or LSND range as discussed above, there can be many resonant oscillation for $\nu_e \leftrightarrow \nu_{\mu,\tau}$, $\nu_e \leftrightarrow \nu_s$ and $\bar{\nu}_{\mu,\tau} \leftrightarrow \bar{\nu}_s$ before they emerge out of the fireball and about 50% of these neutrinos will resonantly convert. These MeV neutrinos signals will be similar to the one from Supernovae, for example SN1987A. Unfortunately with the present generation neutrino telescopes these νs can't be detected due to their cosmological distance and the fluxes are extremely negligible compared to the galactic supernovae.

ACKNOWLEDGMENTS

This work is partially supported by the CONACyT (Mexico) grants No. 32279-E and No. 40025-F.

[1] T. Piran, Phys. Rep. **314**, 575 (1999).
[2] Eli Waxman, Lect. Notes in Phys. **598**, 393 (2003).
[3] T. Piran, Phys. Rep. **333-334**, 529 (2000).
[4] P. Mészáros Nucl. Phys. B (proc. Suppl.) **80**, 63 (2000).
[5] P. Mészáros Annu. Rev. Astron. Astrophys. **40**, 137 (2002).
[6] P. W. Guilbert, A. C. Fabian, and M. J. Rees, Mon. Not. R. Astr. Soc **205**, 593 (1983).
[7] E. V. Derishev, V. V. Kocharovsky, and VI. V Kocharovsky, Astron. Astrophys. **345**, L51 (1999).
[8] E. V. Derishev, V. V. Kocharovsky, and VI. V Kocharovsky, Astrophys. J. **521**, 640 (1999).
[9] W. Kluźniak, Astrophys. J. Lett. **508**, L29 (1998).
[10] Raymond R. Volkas, and Yvonne Y. Y. Wong, Astropart. Phys. **13**, 21 (2000).
[11] B. Paczyński, Astrophys. J. Lett. **308**, L43 (1986).
[12] J. Goodman, Astrophys. J. **308**, L47 (1986).
[13] Eli Waxman, and J. N. Bahcall, Phys. Rev. Lett. **78**, 2292 (1997).
[14] T. J. Weiler, W. A. Simmons, S. Pakvasa, and J. G. Learned, hep-ph/9411432.
[15] M. Ruffert, and H.-Th. Janka, Astron. Astrophys. **344**, 573 (1999).
[16] J. Hjorth, el. al., Nature **423**, 847 (2003).
[17] George G. Raffelt, Astrophys. J. **561**, 890 (2001).
[18] D. Notzöld, and G. Raffelt, Nucl. Phys. B **307**, 924 (1988).
[19] K. Enqvist, K. Kainulaien, and J. Maalampi, Nucl. Phys. B **349**, 754 (1991).
[20] J. C. D'Olivo, Manuel Torres, and J. F. Nieves, Phys. Rev. D **46**, 1172 (1992).
[21] S. Sahu, Phys. Rev. D **61**, 023003 (1999).
[22] A. Athar, J. T. Peltoniemi, and A. Yu. Smirnov, Phys. Rev. D **51**, 6647 (1995).
[23] The SNO Collaboration, Phys. Rev. Lett. **92**, 181301-1 (2004).
[24] The KamLAND Collaboration, hep-ex/0406035.
[25] The Super-Kamiokande Collaboration, Phys. Rev. Lett. **93**, 10180-1 (2004).
[26] E. D. Church, K. Eitel, G. B. Mills, and M. Steidl, Phys. Rev. **D66**, 013001 (2002).